

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07B 53/00, C07C 29/143, 35/27, 45/30, 49/697</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/15299</b> <b>(43) International Publication Date:</b> 8 June 1995 (08.06.95)
<b>(21) International Application Number:</b> PCT/IB94/00263 <b>(22) International Filing Date:</b> 2 September 1994 (02.09.94)  <b>(30) Priority Data:</b> 08/159,156 30 November 1993 (30.11.93) US  <b>(60) Parent Application or Grant</b> (63) Related by Continuation US 08/159,156 (CON) Filed on 30 November 1993 (30.11.93)  <b>(71) Applicant (for all designated States except US):</b> PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US).  <b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> QUALLICH, George, J. [US/US]; 349 Norwich-Westerly Road, North Stonington, CT 06359 (US).  <b>(74) Agents:</b> SPIEGEL, Allen, J. et al.; Pfizer Inc., Patent Dept., 235 East 42nd Street, New York, NY 10017 (US).		<b>(81) Designated States:</b> CA, FI, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> PROCESS FOR PREPARING A CHIRAL TETRALONE  <b>(57) Abstract</b>  A process for preparing the chiral (4S)-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone is disclosed wherein racemic 4-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone is asymmetrically reduced by contacting the racemic tetralone with an asymmetric reagent to produce a mixture of cis and trans alcohols, separating the cis from the trans alcohols, and oxidizing the (4S) enantiomer of the resulting cis and trans alcohols. Also disclosed are novel intermediates used in the synthesis of the above chiral tetralone.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

5 PROCESS FOR PREPARING A CHIRAL TETRALONEBackground of the Invention

The present invention relates to a novel process for asymmetrically reducing racemic 4-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone (hereinafter also referred to as "the tetralone" or "the racemic tetralone") and for preparing chiral (4S)-  
10 (3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone (hereinafter also referred to as "the chiral tetralone"), which has utility as an intermediate in the production of pure cis-(1S)(4S)-N-methyl-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthaleneamine (sertraline). Sertraline is a known antidepressant agent. This invention also relates to novel intermediates in the synthesis of chiral tetralone.

15 Several documents relate to the synthesis of pure racemic N-methyl-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthaleneamine starting with 3,4-dichlorobenzophenone and proceeding via racemic (+)-4-(3,4-dichlorophenyl)-4-butanolic acid and then to (+)-4-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone. See, e.g., U.S. Patent Nos. 4,536,518 (August 20, 1985); 4,556,676 (December 3, 1985);  
20 4,777,288 (October 11, 1988); and 4,839,104 (June 13, 1989); and Journal of Medicinal Chemistry, Vol. 27, No. 11, p. 1508 (1984).

Tetrahedron, Vol. 48, No. 47, pp. 10239-10248 (1992) relates to a process for preparing the (4S)-enantiomer of 4-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone comprising reducing the 4-ketobutanolic acid ester with a carbonyl  
25 reducing agent, as outlined in E. J. Corey et al., Journal of Organic Chemistry, Vol. 53, p. 2861 (1988), to ultimately afford chiral tetralone.

Other asymmetric methods of synthesis have been employed in the art, such as those described by W. M. Whitesides et al., Journal of the American Chemical Society, Vol. 91, No. 17, p. 4871 (1969); K. Mori et al., Synthesis, p. 752 (1982); B. H.  
30 Lipshutz et al., Journal of Organic Chemistry, Vol. 49, p. 3928 (1984); B. H. Lipshutz et al., Journal of the American Chemical Society, Vol. 104, p. 4696 (1982); G. M. Whitesides et al., Journal of the American Chemical Society, Vol. 91, No. 17 (1969); C. R. Johnson et al., Journal of the American Chemical Society, Vol. 95, No. 23, p. 7783 (1973); B. H. Lipshutz et al., Tetrahedron, Vol. 40, No. 24, p. 5005 (1984); and C. R.  
35 Johnson et al., Journal of the American Chemical Society, Vol. 95, No. 23, p. 7777 (1973).

-2-


All of the documents cited herein, including the foregoing, are incorporated herein in their entireties.

### Summary of the Invention

Broadly, the present invention relates to a process for asymmetrically reducing  
 5 racemic 4-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone comprising reacting the racemic 4-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone with an asymmetric ketone reducing agent. The asymmetric ketone reducing agent is preferably a catalytic chiral oxazaborolidine compound.

Preferred chiral oxazaborolidine compounds have the formula:

10



15


wherein:

20  $R^1$  is hydrogen,  $(C_1-C_6)$  alkyl, benzyl, phenyl or phenyl substituted with up to three substituents independently selected from  $(C_1-C_6)$  alkyl,  $(C_1-C_6)$  alkoxy and halo; and

$R^2$  and  $R^3$  are syn and the same and are each phenyl or phenyl substituted with up to three substituents independently selected from  $(C_1-C_6)$  alkyl,  $(C_1-C_6)$  alkoxy and halo.

25 Other preferred chiral oxazaborolidine compounds have the formula:

30



-3-

wherein:

$R^4$  is hydrogen, lower alkyl or aralkyl;

$n$  is 2, 3, or 4, such that the group  $(CH_2)_n$  forms, together with the oxazaborolidine nitrogen and adjacent carbon, a 4-, 5- or 6-membered ring; and


5  $R^5$  and  $R^6$  are phenyl.

Another preferred asymmetric ketone reducing agent comprises either enantiomer of the compound having the formula:



wherein  $lpc$  is isopinocampheyl,  $B$  is boron and  $X$  is halo.


10 The reduction of the racemic tetralone, depending on the asymmetric ketone reducing agent chosen, will yield either cis and trans alcohols having the following formulae:



or cis and trans alcohols having the following formulae:

30

-4-



The enantiomer of the asymmetric reducing agent determines whether (I) and  
 15 (III) or (II) and (IV) is produced.


The present invention also relates to each of the two reduction processes  
 described above (i.e., that which produces compounds (I) and (III) and that which  
 produces compounds (II) and (IV)), further comprising separating, respectively, the cis  
 alcohol (I) from the trans alcohol (III) or the cis alcohol (IV) from the trans alcohol (II)  
 20 and oxidizing, respectively, the resulting cis alcohol (I) or trans alcohol (II) to produce  
 chiral tetralone.

The present invention also relates to a process comprising reacting racemic  
 tetralone with an asymmetric ketone reducing agent to produce compounds having the  
 formulae:


25

30

-5-



15 or compounds having the formulae:




30 said process further comprising the steps of oxidizing the compounds having, respectively, formula (III) or (IV) to produce the 4(R) enantiomer of the tetralone ((4R)-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone), and contacting the resulting 4(R) tetralone with a base to produce racemic tetralone.

The present invention also relates to compounds having the following formulae:


-6-

5




10

15



20


25



30

; and

-7-



The term "halo", as used herein, unless otherwise indicated, includes chloro, fluoro, bromo and iodo.

The term "alkyl", as used herein, unless otherwise indicated, includes saturated  
15 monovalent hydrocarbon radicals having straight, branched or cyclic moieties or combinations thereof.

The term "alkoxy", as used herein, includes O-alkyl groups wherein "alkyl" is defined as above.

The term "aralkyl", as used herein, includes aryl groups, wherein "aryl" is defined  
20 as below, terminating in an alkyl group, as defined above, which is the point of attachment.

The term "aryl", as used herein, means mononuclear aromatic hydrocarbon groups such as phenyl, which can be unsubstituted or substituted in one or more positions, and polynuclear aryl groups such as naphthyl, anthryl, phenanthryl, and so  
25 forth, which can be unsubstituted or substituted with one or more groups.


The term "one or more substituents", as used herein, includes from one to the maximum number of substituents possible based on the number of available bonding sites.

#### Detailed Description of the Invention


30 The processes of this invention for preparing the chiral tetralone (4S)-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone are depicted in the following reaction schemes:

-8-

## Scheme 1



## Scheme 2




-9-

Referring to Scheme 1, racemic 4-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone (tetralone) is asymmetrically reduced by reacting the racemic tetralone with an asymmetric reagent (A) or (B), wherein (A) and (B) are enantiomers. Reduction of racemic tetralone with enantiomer A yields compounds of formulae I and III.

5 Reduction of racemic tetralone with enantiomer B yields compounds of formulae II and IV.

The reduction is performed in a suitable solvent such as tetrahydrofuran, toluene, or an alternative etherial solvent. The reduction is performed at a temperature of from about -20°C to about 50°C, preferably from 20°C to 25°C. The ratio of  
 10 racemic tetralone to asymmetric reagent is from about 1.0:0.025 to about 1.0:1.5. When the asymmetric reagent is a compound of formula (V), (Va) or (VI), then the ratio of racemic tetralone to asymmetric reagent is preferably from about 1.0:0.025 to about 1.0:0.1. When the asymmetric reagent is a compound of formula  $\text{Ipc}_2\text{BX}$ , then the ratio of racemic tetralone to asymmetric reagent is preferably from about 1.0:1.0 to about  
 15 1.0:1.5. The asymmetric reduction of racemic tetralone produces a mixture of cis and trans alcohols of the formulae (I) and (III) or of the formulae II and IV depending upon the chirality of the asymmetric reagent employed. The cis alcohol (I) can be separated from the trans alcohol (III) by methods known in the art, such as chromatography. Similarly, the trans alcohol II can be separated from the cis alcohol IV by methods  
 20 known in the art. In each case, the desired product possesses the chirality desired for sertraline. The (4S) tetralone can be prepared by Jones oxidation, Swern oxidation, Manganese dioxide, pyridium chlorochromate, and pyridium dichromate of the resulting cis alcohol (I) and trans alcohol (II).


Examples of suitable asymmetric reducing reagents include chiral  
 25 oxazaborolidine compounds of the formula:



-10-

wherein  $R^1$  is hydrogen,  $(C_1-C_8)$ alkyl, benzyl, phenyl or phenyl substituted with up to three substituents independently selected from  $(C_1-C_8)$ alkyl,  $(C_1-C_8)$ alkoxy or halo; and  $R^2$  and  $R^3$  are syn and the same and are each phenyl or phenyl substituted with up to three substituents independently selected from  $(C_1-C_8)$ alkyl,  $(C_1-C_8)$ alkoxy or halo groups such as chloro or fluoro. A preferred number of substituents is zero. A preferred group of such compounds is the group of compounds wherein  $R^1$ ,  $R^2$  and  $R^3$  are all unsubstituted phenyl. Especially preferred is the compound wherein  $R^2$  and  $R^3$  are each unsubstituted phenyl and  $R^1$  is methyl. Also, especially preferred is the compound wherein  $R^2$  and  $R^3$  are each phenyl and  $R^1$  is hydrogen.

Suitable asymmetric reagents also include a chiral 1,3,2-oxazaborolidine of the formula:



in which:  $R^4$  is hydrogen, lower alkyl or aralkyl;  $n$  is 2, 3, or 4, such that the group  $(CH_2)_n$  forms, together with the oxazaborolidine nitrogen and adjacent carbon, a 4-, 5- or 6-membered ring; and  $R^5$  and  $R^6$  are phenyl. Aralkyl is as defined above. Preferred alkyl groups of the aralkyl are  $CH_3$ . Preferred aralkyl groups are phenylalkyl groups.

Suitable asymmetric reagents also include a haloborane represented by the formula:  $lpc_2BX$ , wherein  $lpc$  is isopinocampheyl,  $B$  is boron and  $X$  is halo.

Additional suitable asymmetric reagents are disclosed in U.S. Patent No. 5,189,177 issued February 23, 1993; U.S. Patent No. 4,943,635 issued July 24, 1990; U.S. Patent No. 4,772,752 issued September 20, 1988; U.S. Patent Application Serial No. 08/061,895 filed May 14, 1993; International Patent Application PCT/US93/00687, filed February 1, 1993; International Patent Application PCT/US92/05434, filed July 1, 1992; and International Patent Application PCT/US92/05433, filed July 1, 1992.

Referring to Scheme 2, the process for making (4S)-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone may optionally contain one or more additional steps wherein alcohols (III) and/or (IV) are recycled. In this process, the alcohols (III) and/or

(IV) are oxidized to produce 4(R) enantiomer of the tetralone, which is then reacted with a base to produce the racemic tetralone. The oxidation can be done by methods known to those skilled in the art. The racemization reaction is performed at a temperature of from about 0°C to about 100°C, preferably 25°C to 65°C. The 4(R) enantiomer of the tetralone is reacted with a base at a temperature of from about 25°C to about 65°C, preferably 50°C to 80°C. Suitable bases for this reaction include potassium *t*-butoxide, sodium hydroxide, sodium methoxide, and potassium hydroxide. A preferred base is potassium *t*-butoxide.

The (4S)-4-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone final product afforded by the process of this invention is a valuable intermediate that can be used to synthesize the antidepressant agent known as sertraline or cis-(1S)(4S)-N-methyl-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthaleneamine by methods disclosed in the previously discussed prior art. More specifically, (4S)-4-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone is first converted to (4S)-N-[4-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenyldine]methanamine and then finally to the desired cis-(1S)(4S)-N-methyl-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthaleneamine by the known methods of the prior art process, as earlier described in U.S. Patent No. 4,536,518 (August 20, 1985). In the present instance, the optically-active ketone, viz., (4S)-4-(3,4-dichlorophenyl)-1(2H)-naphthalenone, is first reductively aminated to give chiral cis-N-methyl-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthaleneamine and the latter product is then separated by chromatographic means to ultimately yield the desired final medicinal product which is sertraline.

The preparation of other compounds of the present invention not specifically described in the foregoing experimental section can be accomplished using combinations of the reactions described above that will be apparent to those skilled in the art.

In each of the reactions discussed or illustrated in Schemes 1 or 2 above, pressure is not critical unless otherwise indicated. Pressures from about 0.9 atmospheres to about 2 atmospheres are generally acceptable and ambient pressure, i.e., about 1 atmosphere, is preferred as a matter of convenience.

The activity, methods for testing activities, dosages, dosage forms, methods of administration and background information concerning sertraline are set forth in U.S. Patent Nos. 4,536,518 (August 20, 1985), 4,777,288 (October 11, 1988), and 4,839,104

(June 13, 1989), and the Journal of Medicinal Chemistry, Vol. 27, No. 11, p. 1508 (1984).

The present invention is illustrated by the following examples. It will be understood, however, that the invention is not limited to the specific details of these  
5 examples.

#### EXAMPLE 1

##### (4S)-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone

Borane methylsulfide complex (2M in THF, 6.0 mL, 12 mmol) was added all at once to a solution of (1S, 2R)-(+)-erythro-2-amino-1,2-diphenylethanol [J. Amer. Chem. Soc. 1216 (1951) also commercially available] (183 mg, 0.86 mmol) in THF (55 mL)  
10 under a nitrogen atmosphere. The solution was stirred for 18 hours. Racemic tetralone (5.0 g, 17 mmol) as a solution in THF was added over 1 hour, the reaction stirred 15 minutes after the addition was completed, cooled to 0°C and quenched with methanol. After stirring the quenched reaction for 18 hours the solvents were removed under  
15 vacuum, the contents dissolved in methylene chloride (100 mL), and washed sequentially with pH4 phosphate buffer (100 mL), water (100 mL), treated with magnesium sulfate, and solvent removed to afford a mixture of the cis and trans alcohols (5.01 g). Chromatography with ethyl acetate/hexanes provided the less polar cis alcohol. <sup>1</sup>H NMR δ (CDCl<sub>3</sub>) 7.46 (dd, J=1Hz, J=7Hz, 1H), 7.41-7.07 (m, 4H), 6.98  
20 (dd, J=2Hz, J=8Hz, 1H), 6.82 (d, J=7Hz, 1H), 4.86 (t, J=4Hz, 1H), 3.99 (t, J=8Hz, 1H), 2.18-1.87 (m, 5H). <sup>13</sup>C NMR δ 147.0, 138.9, 138.4, 132.4, 130.7, 130.4, 130.2, 129.8, 129.1, 128.3, 128.2, 127.1, 67.9, 45.1, 30.1, 28.2 and the more polar trans alcohol. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.54 (d, J=7Hz, 1H), 7.4-7.07 (m, 4H), 6.90-6.75 (m, 2H), 4.88 (t, J=5Hz, 1H), 4.13 (t, J=6Hz, 1H), 2.43-1.63 (m, 5H). The less polar cis alcohol 1 (160  
25 mg, 0.546 mmol) was dissolved in methylene chloride (5 mL), treated with pyridium chlorochromate (220 mg, 1.023 mmol), and stirred for 2 hours at ambient temperature. Diethyl ether was added (25 mL), stirred 20 minutes, and the solvent decanted. The residual dark gum was washed with diethyl ether (2 X 15 mL), the organic layers combined, filtered through a pad of magnesium sulfate, and solvent removed under  
30 vacuum to afford a brown oil (170 mg). Chromatography on silica (5.1 g) eluting with methylene chloride provided the chiral tetralone as a clear oil (118 mg). This material was determined to be ≥ 95% ee by HPLC with a chiral support (Diacel Co. ChiralcelOD 4.6 mm X 25 cm, 10% isopropyl alcohol/hexane).

-13-

EXAMPLE 2(4S)-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone

(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone ("the tetralone") (5 g, 17 mmol), (S)-Tetrahydro-1-methyl-3,3-diphenyl-1H,3H-pyrrolo-[1,2-c][1,3,2]oxazaborole [J. Org. Chem. 2861 (1988)] (238 mg, 0.859 mmol), and tetrahydrofuran (THF) (68 mL) were combined at ambient temperature in a flamed dried flask under a nitrogen atmosphere. Borane methylsulfide complex (2 M in THF, 4.56 mL) was added over 1 hour, 30 minutes later the reaction was quenched at 0°C with methanol (15.8 mL), and stirred for 18 hours. The solvents were removed under vacuum, the contents dissolved in methylene chloride (68 mL), and washed sequentially with pH4 phosphate buffer (68 mL), water (68 mL), treated with magnesium sulfate, and solvent removed to afford a mixture of the cis and trans alcohols (4.93 g). Chromatography with ethylacetate/hexanes provided the less polar cis alcohol  $\alpha_D = -52.27$  ( $c = 1.01$ , methylene chloride) and more polar trans alcohol  $\alpha_D = +37.79$  ( $c = 1.18$ , methylene chloride). The more polar trans alcohol 2 (160 mg, 0.546 mmol) was dissolved in methylene chloride (5 mL), treated with pyridium chlorochromate (220 mg, 1.023 mmol), and stirred for 2 hours at ambient temperature. Diethyl ether was added (25 mL), stirred 20 minutes, and the solvent decanted. The residual dark gum was washed with diethyl ether (2 X 15 mL), the organic layers combined, filtered through a pad of magnesium sulfate, and solvent removed under vacuum to afford a brown oil (162 mg). Chromatography on silica (5 g) eluting with 25% ethyl acetate/hexanes provided the chiral tetralone as a clear oil (139 mg)  $\alpha_D = +36.8$  ( $c = 1.11$ ) which corresponds to a 56% ee.

EXAMPLE 3(4S)-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone

[(+)- $\beta$ -chlorodiisopinocampheylborane] (6.07 g, 18.97 mmol) was dissolved in THF (13.6 mL) under a nitrogen atmosphere, and cooled to -25°C. Tetralone (5 g, 17 mmol) was added as a solution in THF (13.6 mL), the contents allowed to warm to ambient temperature, and stirred 46 hours. The solvent was removed under vacuum, diethyl ether (65 mL) and ethanol amine (3.9 mL) were added, and the contents stirred for 18 hours. The precipitate was filtered off, washed with pentane (2 X 20 mL), and the solvent removed under vacuum from the filtrate to yield the crude product (4.98 g) which was chromatographed on silica with 25% ethyl acetate/hexanes to separate the cis and trans alcohols. The more polar trans alcohol 2 (175 mg, 0.579 mmol) was

-14-

dissolved in methylene chloride (5 mL) and treated with pyridium chlorochromate (192 mg) for 2 hours at ambient temperature. Diethyl ether was added (25 mL), stirred 15 minutes, and the solvents decanted. The residual black semisolid was washed with diethyl ether (2 X 10 mL), the organic phases combined, filtered through CELITE, and solvent removed under vacuum to yield the crude chiral tetralone. Chromatography on silica eluting with 25% ethyl acetate/hexanes afforded 165 mg of pure product  $\alpha_D = +30.94$  ( $c = 1.28$ , acetone) which corresponds to 47% ee.

#### EXAMPLE 4


##### 4-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone

Pyridium chlorochromate oxidation of the trans alcohols 3 and 4 with the same procedure employed on alcohols 1 and 2 provided the chiral tetralone. Racemization of the chiral tetralone into racemic tetralone was achieved as follows. Potassium t-butoxide (90 mg, 0.80 mmol) was added to a solution of chiral tetralone (1.12 g, 3.84 mmol) in THF (4 mL). The solution was refluxed for 18 hours under nitrogen, cooled to ambient temperature, methylene chloride (10 mL) and aqueous hydrochloric acid (1N, 20 mL) added, and the phases separated. The organic phase was washed with water (10 mL), brine (10 mL), dried with magnesium sulfate, and solvent removed under vacuum to yield 1.1 g of the crude racemic tetralone. Recrystallization from methanol afforded 1.07 g (95%) of the racemic tetralone mp-104-5°C. Other base solvent combinations which effect racemization are methanol/sodium methoxide, methanol/sodium hydroxide, and methanol/potassium hydroxide.

-15-

CLAIMS


1. A process for asymmetrically reducing racemic 4-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone comprising reacting the racemic 4-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone with an asymmetric ketone reducing agent.
2. A process according to claim 1 wherein the asymmetric ketone reducing agent is a chiral catalytic oxazaborolidine compound.
3. A process according to claim 2 wherein the chiral oxazaborolidine compound has the formula:




wherein:

- $R^1$  is hydrogen,  $(C_1-C_6)$  alkyl, benzyl, phenyl or phenyl substituted with up to three substituents independently selected from  $(C_1-C_6)$  alkyl,  $(C_1-C_6)$  alkoxy and halo; and
- $R^2$  and  $R^3$  are syn and the same and are each phenyl or phenyl substituted with up to three substituents independently selected from  $(C_1-C_6)$  alkyl,  $(C_1-C_6)$  alkoxy and halo.

4. A process according to claim 2 wherein the chiral oxazaborolidine compound has the formula:



-17-



8. A process according to claim 6 wherein said process further comprises  
15 the steps of separating the cis alcohol (I) from the trans alcohol (III) and oxidizing the  
the resulting cis alcohol (I) to produce chiral tetralone.

9. A process according to claim 7 wherein said process further comprises  
the steps of separating the cis alcohol (IV) from the trans alcohol (II) and oxidizing the  
the resulting trans alcohol (II) to produce chiral tetralone.

20 10. A process according to claim 1 wherein the reaction of racemic tetralone  
with an asymmetric ketone reducing agent yields compounds having the formulae:

25

30

-16-

wherein:

$R^4$  is hydrogen, lower alkyl or aralkyl;

$n$  is 2, 3, or 4, such that the group  $(CH_2)_n$  forms, together with the oxazaborolidine nitrogen and adjacent carbon, a 4-, 5- or 6-membered ring; and


5  $R^5$  and  $R^6$  are phenyl.

5. A process according to claim 1 wherein the asymmetric ketone reducing agent comprises either enantiomer of the compound having the formula:



wherein  $lpc$  is Isopinocampheyl,  $B$  is boron and  $X$  is halo.

10 6. A process according to claim 1 wherein the reduction of the racemic tetralone yields cis and trans alcohols having the following formulae:




25


7. A process according to claim 1 wherein the reduction of the racemic tetralone yields cis and trans alcohols having the following formulae:

30

-18-




15 or compounds having the formulae:



11. A process according to claim 10, further comprising the steps of  
30 oxidizing the compounds having formula (III) or (IV) to produce (4R)-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone, and contacting the resulting 4(R) tetralone with a base to produce racemic 4-(3,4-dichlorophenyl)-3,4-dihydro-1(2H)-naphthalenone.

-19-

12. A process according to claim 10 wherein said asymmetric ketone reducing agent is a chiral oxazaborolidine having the formula:




wherein:

R<sup>1</sup> is hydrogen, (C<sub>1</sub>-C<sub>8</sub>) alkyl, benzyl, phenyl or phenyl substituted with up to three substituents independently selected from (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>1</sub>-C<sub>8</sub>)alkoxy and halo; and

15 R<sup>2</sup> and R<sup>3</sup> are syn and the same and are each phenyl or phenyl substituted with up to three substituents independently selected from (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>1</sub>-C<sub>8</sub>)alkoxy and halo.

13. A process according to claim 10 wherein said asymmetric ketone reducing agent is a chiral oxazaborolidine having the formula:



wherein:


R<sup>4</sup> is hydrogen, lower alkyl or aralkyl;

n is 2, 3, or 4, such that the group (CH<sub>2</sub>)<sub>n</sub> forms, together with the oxazaborolidine nitrogen and adjacent carbon, a 4-, 5- or 6-membered ring; and


30 R<sup>5</sup> and R<sup>6</sup> are phenyl.

-20-

14. A compound of the formula:




15. A compound of the formula:




-21-

16. A compound of the formula:



17. A compound of the formula:



## INTERNATIONAL SEARCH REPORT

Int. l. Application No.

PCT/IB 94/00263

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07B53/00 C07C29/143 C07C35/27 C07C45/30 C07C49/697

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO,A,93 23408 (PFIZER) 25 November 1993 see claims; examples 4,12 ---	1-3
A	EP,A,0 305 180 (PRESIDENT AND FELLOWS OF HARVARD UNIVERSITY) 1 March 1989 see page 18, line 44; claims; example 2 ---	1,2,4
A	US,A,4 772 752 (H BROWN) 20 September 1988 see claims; examples -----	1,5



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"B" document member of the same patent family

Date of the actual completion of the international search

17 October 1994

Date of mailing of the international search report

- 8. 11. 94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patendaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2046, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Heywood, C

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 94/00263

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9323408	25-11-93	AU-B- 3593193 HU-A- 65136	13-12-93 28-04-94
EP-A-0305180	01-03-89	US-A- 4943635	24-07-90
US-A-4772752	20-09-88	US-A- 4866181 US-A- 5043479	12-09-89 27-08-91